

Opposite-sign Dilepton Asymmetry of Neutral B Decays: a Probe of New Physics from CPT or $\Delta B = \Delta Q$ Violation

Zhi-zhong Xing¹

Sektion Physik, Universität München, Theresienstrasse 37A, 80333 München, Germany

Abstract

We show that new physics, either from CPT violation in B^0 - $\bar{B}^0$ mixing or from $\Delta B = -\Delta Q$ transitions, may lead to an opposite-sign dilepton asymmetry of neutral B -meson decays. Both effects have the same time-dependent behavior and therefore are in general indistinguishable from each other. We also clarify some ambiguity associated with the constraint on CPT violation in semileptonic decays of incoherent B^0 and $\bar{B}^0$ mesons.

¹ E-mail: Xing@hep.physik.uni-muenchen.de

The B -meson factories under construction are going to provide a unique opportunity for the study of CP violation, both its phenomenon and its origin. At the second or final stage of these facilities sufficient $B\bar{B}$ events (e.g., $\mathcal{N}_{B\bar{B}} \geq 10^9$) will be accumulated, then a direct test of the CPT symmetry and the $\Delta B = \Delta Q$ rule should be experimentally feasible. This is expected to open a new window for probing new physics beyond the standard model, as both CPT and $\Delta B = \Delta Q$ conservation laws work extremely well within the standard model. Up to now some phenomenological analyses of possible CPT -violating signals in semileptonic and nonleptonic decays of neutral B mesons have been made in Refs. [1] – [6], whose results depend on the validity of the $\Delta B = \Delta Q$ rule. The possibility to detect the effect of $\Delta B = -\Delta Q$ transitions at B factories has been investigated in Ref. [7] in the assumption of CPT invariance.

It was noticed in Refs. [1] – [6] that CPT violation in B^0 - $\bar{B}^0$ mixing could lead to an opposite-sign dilepton asymmetry of neutral- B decays, or more generally, an asymmetry between $B^0(t) \rightarrow B^0 \rightarrow l^+ X^-$ and $\bar{B}^0(t) \rightarrow \bar{B}^0 \rightarrow l^- X^+$ decay rates. Applying this idea to the time-dependent semileptonic decays of B^0 and $\bar{B}^0$ mesons at the Z resonance, the OPAL Collaboration obtained a constraint on the CPT -violating parameter δ_B , i.e., $\text{Im}\delta_B = -0.020 \pm 0.016 \pm 0.006$ [8]. The ideal experimental environment for measuring the time distribution of opposite-sign dilepton events will be at the KEK and SLAC asymmetric B -meson factories, where $B^0\bar{B}^0$ pairs can coherently be produced through the decay of the $\Upsilon(4S)$ resonance.

Allowing both CPT violation and $\Delta B = -\Delta Q$ transitions, we shall carry out a new analysis of the opposite-sign dilepton asymmetry of neutral B decays and clarify some ambiguity associated with the constraint on CPT violation in semileptonic decays of incoherent B^0 and $\bar{B}^0$ mesons. It is shown that the effect of $\Delta B = -\Delta Q$ transitions and that of CPT violation have the same time-dependent behavior in the opposite-sign dilepton events, therefore they are in general indistinguishable from each other. We point out that a meaningful constraint on $\text{Im}\delta_B$ depends actually on the smallness of the decay width difference between two neutral- B mass eigenstates, on the condition $|\text{Re}\delta_B| \leq |\text{Im}\delta_B|$, and on the validity of the $\Delta B = \Delta Q$ rule.

Mixing between B^0 and $\bar{B}^0$ mesons arises naturally from their common coupling to a subset of real and virtual intermediate states, hence the mass eigenstates $|B_1\rangle$ and $|B_2\rangle$ are different from the flavor eigenstates $|B^0\rangle$ and $|\bar{B}^0\rangle$. A *non-linear* parametrization of CP - and CPT -violating effects in B^0 - $\bar{B}^0$ mixing reads as

$$\begin{aligned} |B_1\rangle &= \cos\frac{\theta}{2} e^{-i\frac{\phi}{2}} |B^0\rangle + \sin\frac{\theta}{2} e^{+i\frac{\phi}{2}} |\bar{B}^0\rangle, \\ |B_2\rangle &= \sin\frac{\theta}{2} e^{-i\frac{\phi}{2}} |B^0\rangle - \cos\frac{\theta}{2} e^{+i\frac{\phi}{2}} |\bar{B}^0\rangle, \end{aligned} \quad (1)$$

where θ and ϕ are in general complex, and the normalization factors of $|B_1\rangle$ and $|B_2\rangle$ have

been neglected. CPT invariance requires $\cos\theta = 0$, while CP conservation requires both $\cos\theta = 0$ and $\phi = 0$ [9]. In some literature the *linear* parametrization ²

$$\begin{aligned} |B_1\rangle &= (1 + \epsilon_B + \delta_B)|B^0\rangle + (1 - \epsilon_B - \delta_B)|\bar{B}^0\rangle, \\ |B_2\rangle &= (1 + \epsilon_B - \delta_B)|B^0\rangle - (1 - \epsilon_B + \delta_B)|\bar{B}^0\rangle, \end{aligned} \quad (2)$$

together with the conventions $|\epsilon_B| \ll 1$ and $|\delta_B| \ll 1$, has been adopted (here again the normalization factors of $|B_1\rangle$ and $|B_2\rangle$ are neglected). It is straightforward to find the relationship between (θ, ϕ) and (δ_B, ϵ_B) parameters in the leading-order approximation, $\epsilon_B \approx i\phi/2$ and $\delta_B \approx \cot\theta/2$, provided the conventions $|\cos\theta| \ll 1$ and $|\phi| \ll 1$ are taken. Note that the requirement $|\epsilon_B| \ll 1$ (or $|\phi| \ll 1$) corresponds to an exotic phase convention of quark fields, which is incompatible with several popular parametrizations of the quark flavor mixing matrix [10]. In subsequent calculations we shall mainly make use of the (θ, ϕ) notation (in which ϕ may take arbitrary values), and translate it into the (δ_B, ϵ_B) notation when necessary.

The proper-time evolution of an initially pure $|B^0\rangle$ or $|\bar{B}^0\rangle$ state is then given as [2]

$$\begin{aligned} |B^0(t)\rangle &= e^{-(im + \frac{\Gamma}{2})t} \left[g_+(t)|B^0\rangle + \tilde{g}_+(t)|\bar{B}^0\rangle \right], \\ |\bar{B}^0(t)\rangle &= e^{-(im + \frac{\Gamma}{2})t} \left[g_-(t)|\bar{B}^0\rangle + \tilde{g}_-(t)|B^0\rangle \right], \end{aligned} \quad (3)$$

where

$$\begin{aligned} g_{\pm}(t) &= \cosh\left(\frac{ix - y}{2}\Gamma t\right) \pm \cos\theta \sinh\left(\frac{ix - y}{2}\Gamma t\right), \\ \tilde{g}_{\pm}(t) &= \sin\theta e^{\pm i\phi} \sinh\left(\frac{ix - y}{2}\Gamma t\right). \end{aligned} \quad (4)$$

In Eqs. (3) and (4) we have defined $m \equiv (m_1 + m_2)/2$, $\Gamma \equiv (\Gamma_1 + \Gamma_2)/2$, $x \equiv (m_2 - m_1)/\Gamma$ and $y \equiv (\Gamma_1 - \Gamma_2)/(2\Gamma)$, where $m_{1,2}$ and $\Gamma_{1,2}$ are the mass and width of $B_{1,2}$. The mixing parameter $x \approx 0.7$ has been measured [11], and $y \sim O(10^{-2})$ is theoretically expected [12]. To calculate the time distribution of opposite-sign dilepton events on the $\Upsilon(4S)$ resonance, we neglect possible tiny effects from electromagnetic final-state interactions and assume CPT invariance in the direct transition amplitudes of semileptonic B decays. Such an assumption can be examined, without the mixing-induced complexity, by measuring the charge asymmetry of semileptonic $B^{\pm}$ decays. The effect of possible $\Delta B = -\Delta Q$ transitions need be taken into account. We then write the relevant semileptonic decay amplitudes as follows:

$$\begin{aligned} \langle l^+ | B^0 \rangle &= A_l, & \langle l^+ | \bar{B}^0 \rangle &= \sigma_l A_l; \\ \langle l^- | \bar{B}^0 \rangle &= A_l^*, & \langle l^- | B^0 \rangle &= \sigma_l^* A_l^*, \end{aligned} \quad (5)$$

²Here we use the notation (δ_B, ϵ_B) instead of (δ, ϵ) , as the latter is commonly adopted to describe CPT - and CP -violating effects in the K^0 - $\bar{K}^0$ mixing system.

where σ_l measures the $\Delta B = -\Delta Q$ effect and $|\sigma_l| \ll 1$ holds.

$|\sigma_l| \neq 0$ implies that it is in practice impossible to have a *pure* tagging of the B^0 or $\bar{B}^0$ state through its semileptonic decay (to l^+ or l^-). For $B^0\bar{B}^0$ pairs produced *incoherently* on the Z resonance or elsewhere, the assumption $|\sigma_l| = 0$ (i.e., the exact $\Delta B = \Delta Q$ rule) is indeed a necessary condition to get pure flavor tagging of B^0 and $\bar{B}^0$ mesons and to study possible tiny violation of CPT symmetry in their semileptonic decays³.

The pure flavor tagging is however unnecessary for the study of fine effects induced by CPT violation and (or) $\Delta B = -\Delta Q$ transitions in opposite-sign dilepton products of *coherent* $B^0\bar{B}^0$ pairs on the $\Upsilon(4S)$ resonance. Here the decay of one neutral B meson at proper time t_1 into a semileptonic state (e.g., $e^\pm X_e^\mp$) may only serve as a rough flavor tagging for the other meson decaying at proper time t_2 into another semileptonic state (e.g., $\mu^\mp X_\mu^\pm$). The overall final state is therefore an opposite-sign dilepton event. Since we are only interested in how the decay rate depends on the time difference $t_2 - t_1$ at an asymmetric B factory, we just take $t_1 = 0$ and $t_2 = t$ with the convention $t > 0$ for the wave function of the coherent $B^0\bar{B}^0$ pair:

$$\Psi(t) = \frac{1}{\sqrt{2}} \left[|B^0(0)\rangle \otimes |\bar{B}^0(t)\rangle - |B^0(t)\rangle \otimes |\bar{B}^0(0)\rangle \right]. \quad (6)$$

Then the rates of two opposite-sign dilepton decays (with $l^\pm$ events at $t = 0$) read

$$\begin{aligned} \mathcal{R}(t) &\equiv |\langle l^+ l'^- | \Psi(t) \rangle|^2 = \frac{1}{2} |A_l|^2 |A_{l'}|^2 e^{-\Gamma t} \left| g_-(t) + \tilde{g}_-(t) \sigma_{l'}^* - \tilde{g}_+(t) \sigma_l \right|^2, \\ \bar{\mathcal{R}}(t) &\equiv |\langle l^- l'^+ | \Psi(t) \rangle|^2 = \frac{1}{2} |A_l|^2 |A_{l'}|^2 e^{-\Gamma t} \left| g_+(t) + \tilde{g}_+(t) \sigma_{l'} - \tilde{g}_-(t) \sigma_l^* \right|^2, \end{aligned} \quad (7)$$

where only the contributions of $O(|\sigma_l|)$ and $O(|\sigma_{l'}|)$ are kept, and the functions $g_\pm(t)$ and $\tilde{g}_\pm(t)$ can be found in Eq. (3). We arrive finally at

$$\begin{aligned} \mathcal{R}(t) &= \frac{1}{4} |A_l|^2 |A_{l'}|^2 e^{-\Gamma t} \left[\cosh(y\Gamma t) + 2\text{Re}\Omega \sinh(y\Gamma t) \right. \\ &\quad \left. + \cos(x\Gamma t) + 2\text{Im}\Omega \sin(x\Gamma t) \right], \\ \bar{\mathcal{R}}(t) &= \frac{1}{4} |A_l|^2 |A_{l'}|^2 e^{-\Gamma t} \left[\cosh(y\Gamma t) - 2\text{Re}\bar{\Omega} \sinh(y\Gamma t) \right. \\ &\quad \left. + \cos(x\Gamma t) - 2\text{Im}\bar{\Omega} \sin(x\Gamma t) \right], \end{aligned} \quad (8)$$

where

$$\begin{aligned} \Omega &= \cos\theta + \xi_{ll'} \sin\theta, \\ \bar{\Omega} &= \cos\theta + \bar{\xi}_{ll'} \sin\theta, \end{aligned} \quad (9)$$

³Note that the treatment in Ref. [3], which takes $|\sigma_l| \neq 0$ on the one hand but assumes the pure flavor tagging of B^0 and $\bar{B}^0$ states on the other hand, is controversial in the experimental environment where $B^0\bar{B}^0$ pairs are incoherently produced.

and

$$\begin{aligned}\xi_{ll'} &= \sigma_l e^{+i\phi} - \sigma_{l'}^* e^{-i\phi}, \\ \bar{\xi}_{ll'} &= \sigma_{l'} e^{+i\phi} - \sigma_l^* e^{-i\phi}.\end{aligned}\tag{10}$$

In obtaining Eq. (8) we have neglected the contributions of $O(|\Omega|^2)$ and $O(|\bar{\Omega}|^2)$.

Obviously the $\Delta B = -\Delta Q$ effects are signified by the rephasing-invariant parameters $\xi_{ll'}$ and $\bar{\xi}_{ll'}$. $\xi_{ll'} = \bar{\xi}_{ll'}$ holds for $l' = l$, and $\xi_{ll'} = -\bar{\xi}_{ll'}^*$ holds if $\text{Im}\phi = 0$. We find that $\xi_{ll'}$ or $\bar{\xi}_{ll'}$ has the same time-dependent behavior as the CPT -violating parameter $\cos\theta$ in the decay rate $\mathcal{R}(t)$ or $\bar{\mathcal{R}}(t)$. *This important feature implies that it is in general impossible to distinguish between the effects of CPT violation and $\Delta B = -\Delta Q$ transitions in the opposite-sign dilepton events, unless one of them is remarkably smaller than the other.*

For illustration we simplify the result in Eq. (8) by taking $\sigma_l = \sigma_{l'} = 0$, i.e, no $\Delta B = -\Delta Q$ transition involved. In this case one gets

$$\begin{aligned}\text{Re}(\Omega + \bar{\Omega}) &= 2\text{Re}(\cos\theta), \\ \text{Im}(\Omega + \bar{\Omega}) &= 2\text{Im}(\cos\theta),\end{aligned}\tag{11}$$

which results in an asymmetry between the decay rates $\mathcal{R}(t)$ and $\bar{\mathcal{R}}(t)$. Defining $\mathcal{A}(t)$ as the ratio of $\mathcal{R}(t) - \bar{\mathcal{R}}(t)$ to $\mathcal{R}(t) + \bar{\mathcal{R}}(t)$, we then have

$$\mathcal{A}(t) = 2 \frac{\text{Re}(\cos\theta) \sinh(y\Gamma t) + \text{Im}(\cos\theta) \sin(x\Gamma t)}{\cosh(y\Gamma t) + \cos(x\Gamma t)},\tag{12}$$

which is independent of the CP -violating parameter ϕ . This formula has already been obtained in Refs. [4] and [6]⁴. It should be noted that the asymmetry $\mathcal{A}(t)$, due to its time dependence, is not restricted to the range $[-1, +1]$. In the limit $y = 0$, Eq. (12) turns out to be $\mathcal{A}(t) = 2\text{Im}(\cos\theta) \tan(x\Gamma t/2)$, which becomes infinity on the point $\Gamma t = \pi/x$ (i.e., around $\Gamma t \approx 4.5$).

If one translates $\cos\theta$ into δ_B , then $\text{Re}(\cos\theta) \approx 2\text{Re}\delta_B$ and $\text{Im}(\cos\theta) \approx 2\text{Im}\delta_B$. Therefore a constraint on $\text{Im}\delta_B$ is achievable from Eq. (12) with $y = 0$. This is indeed the case taken by the OPAL Collaboration in their measurement [8], where the rate difference between $B^0(t) \rightarrow B^0 \rightarrow l^+ X^-$ and $\bar{B}^0(t) \rightarrow \bar{B}^0 \rightarrow l^- X^+$ transitions is essentially equivalent to the asymmetry of opposite-sign dilepton events of $B^0 \bar{B}^0$ decays under discussion. As the theoretical calculation favors $y/x \sim 10^{-2}$ [12], the approximation $y \approx 0$ made in Ref. [8] seems quit reasonable.

However, such an approximation may not be valid if $|\text{Re}\delta_B| \gg |\text{Im}\delta_B|$ holds. Although there is no experimental information about the relative magnitude of $\text{Re}\delta_B$ and $\text{Im}\delta_B$, a

⁴Note that the definitions of y in Refs. [4] and [6] are different in sign. Taking this into account, one can get full consistency between the results in these two papers.

string theory with spontaneous CPT violation and Lorentz symmetry breaking predicts [3]

$$r \equiv \frac{\text{Re}\delta_B}{\text{Im}\delta_B} = \pm \frac{x}{y} \sim \pm 10^2. \quad (13)$$

In this case, the $\sinh(y\Gamma t)$ and $\sin(x\Gamma t)$ terms in the numerator of $\mathcal{A}(t)$ are comparable in magnitude even for small t . Then a meaningful constraint on $\text{Im}\delta_B$ becomes impossible. To illustrate, we plot the time distribution of $\mathcal{A}(t)$ in Fig. 1 with the inputs $r = \pm 1$ and $r = \pm 100$ (for fixed $\text{Im}\delta_B$), respectively. It is obvious in the latter case that the $\text{Re}\delta_B$ contribution to $\mathcal{A}(t)$ is significant (even dominant for large Γt) and a separate bound on $\text{Im}\delta_B$ cannot be obtained.

Next let us simplify the result in Eq. (8) by taking $\cos\theta = 0$ or $\sin\theta = 1$, i.e., no CPT violation in B^0 - $\bar{B}^0$ mixing involved. In this case we obtain

$$\begin{aligned} \text{Re}(\Omega + \bar{\Omega}) &= 2 \sinh(\text{Im}\phi) [\text{Im}Z_{ll'} \sin(\text{Re}\phi) - \text{Re}Z_{ll'} \cos(\text{Re}\phi)] , \\ \text{Im}(\Omega + \bar{\Omega}) &= 2 \cosh(\text{Im}\phi) [\text{Im}Z_{ll'} \cos(\text{Re}\phi) + \text{Re}Z_{ll'} \sin(\text{Re}\phi)] , \end{aligned} \quad (14)$$

where $Z_{ll'} \equiv \sigma_l + \sigma_{l'}$ with $|Z_{ll'}| \ll 1$. Note that $\text{Im}\phi$ describes the CP -violating effect induced by B^0 - $\bar{B}^0$ mixing and can in principle be measured (or constrained) from the same-sign dilepton asymmetry of neutral- B decays (see, e.g., Refs. [1, 4]). The magnitude of $\text{Im}\phi$ is expected to be of $O(10^{-3})$ within the standard model, but it might be enhanced to $O(10^{-2})$ if there exists the CP -violating new physics in B^0 - $\bar{B}^0$ mixing [13]. Anyway the $\text{Re}(\Omega + \bar{\Omega})$ term is doubly suppressed by $\text{Im}\phi$ and $|Z_{ll'}|$, therefore it can be neglected in the rate difference between $\mathcal{R}(t)$ and $\bar{\mathcal{R}}(t)$. The asymmetry $\mathcal{A}'(t)$, defined as the ratio of $\mathcal{R}(t) - \bar{\mathcal{R}}(t)$ to $\mathcal{R}(t) + \bar{\mathcal{R}}(t)$, is then governed by the $\text{Im}(\Omega + \bar{\Omega})$ term with $\cosh(\text{Im}\phi) \approx 1$ as follows:

$$\mathcal{A}'(t) = 2 \frac{[\text{Im}Z_{ll'} \cos(\text{Re}\phi) + \text{Re}Z_{ll'} \sin(\text{Re}\phi)] \sin(x\Gamma t)}{\cosh(y\Gamma t) + \cos(x\Gamma t)}. \quad (15)$$

This instructive result implies that a signal of $\Delta B = -\Delta Q$ transitions can emerge from the opposite-sign dilepton asymmetry of neutral- B decays, if CPT violation is absent or negligibly small. Whether the effect of $\Delta B = \Delta Q$ violation is significant or not beyond the standard model, however, remains an open question.

Comparing Eq. (15) with Eq. (12), one can see that the neglect of the $\sinh(y\Gamma t)$ term in $\mathcal{A}'(t)$ is much safer than that in $\mathcal{A}(t)$. In general, the $\text{Re}(\Omega + \bar{\Omega})$ and $\text{Im}(\Omega + \bar{\Omega})$ terms may consist of comparable effects from CPT violation and $\Delta B = -\Delta Q$ transitions. Hence a constraint on $\text{Im}(\cos\theta)$ or $\text{Im}\delta_B$, due to the presence of $\Delta B = -\Delta Q$ contamination, becomes impossible in the opposite-sign dilepton events.

Keeping with the current experimental interest in tests of discrete symmetries and conservation laws at the upcoming B factories, we have made a new analysis of the opposite-sign

dilepton decays of neutral- B mesons. It is shown that the effects of $\Delta B = -\Delta Q$ transitions and CPT violation have the same time-dependent behavior in such dilepton events, hence they are in general indistinguishable from each other. To separate one kind of new physics from the other will require some delicate measurements of the opposite-sign dilepton events, the same-sign dilepton events, and other relevant nonleptonic decay channels. We remark that an experimental bound on the CPT -violating parameter $\text{Im}\delta_B$ relies on the assumptions $y \approx 0$ and $|\text{Re}\delta_B| \leq |\text{Im}\delta_B|$ as well as the validity of the $\Delta B = \Delta Q$ rule. For this reason the value obtained in Ref. [8] can only shed limited light on CPT invariance or its possible violation in the B^0 - $\bar{B}^0$ mixing system. However, a new analysis of the old data is likely to give a constraint on the imaginary part of $(\Omega + \bar{\Omega})$, provided its real part is not significant.

We hope that a useful test of CPT and $\Delta B = \Delta Q$ conservation laws can finally be realized at the asymmetric B -meson factories. The prospect of such ambitious experiments, as discussed in Ref. [6], is not dim. And the relevant theoretical motivation seems encouraging too [14]. On the phenomenological side, further and more general analyses of CP -violating, CPT -violating and $\Delta B = -\Delta Q$ effects on all types of neutral B -meson decays are therefore desirable.

The author likes to thank M. Jimack for his helpful comments on this work. He is also grateful to V.A. Kostelecký, L.B. Okun, Y. Sakai and S.Y. Tsai for useful discussions.

References

- [1] M. Kobayashi and A.I. Sanda, Phys. Rev. Lett. **69**, 3139 (1992).
- [2] Z.Z. Xing, Phys. Rev. D **50**, 2957 (1994).
- [3] D. Colladay and V.A. Kostelecký, Phys. Lett. B **344**, 259 (1995); V.A. Kostelecký and R. Van Kooten, Phys. Rev. D **54**, 5585 (1996).
- [4] Z.Z. Xing, Phys. Rev. D **55**, 196 (1997). The generic results obtained in Appendix B of this paper are valid for both neutral- D and neutral- B decays.
- [5] P. Colangelo and G. Corcella, Eur. Phys. J. C **1**, 515 (1998).
- [6] A. Mohapatra, M. Satpathy, K. Abe, and Y. Sakai, Phys. Rev. D **58**, 036003 (1998).
- [7] G.V. Dass and K.V.L. Sarma, Phys. Rev. Lett. **72**, 191 (1994).
- [8] OPAL Collaboration, R. Ackerstaff *et al.*, Z. Phys. C **76**, 401 (1997); see also, DELPHI Collaboration, M. Feindt *et al.*, Report No. DELPHI 97-98 CONF 80 (July 1997).
- [9] T.D. Lee and C.S. Wu, Annu. Rev. Nucl. Sci. **16**, 471 (1966).
- [10] We thank Y. Nir for useful discussions. See also, Z.Z. Xing, hep-ex/9709005. For a classification of parametrizations of the quark flavor mixing matrix, see: H. Fritzsch and Z.Z. Xing, Phys. Rev. D **57**, 594 (1998); Phys. Lett. B **413**, 396 (1997).
- [11] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. **3**, 1 (1998).
- [12] I.I. Bigi, V.A. Khoze, N.G. Uraltsev, and A.I. Sanda, in *CP Violation*, edited by C. Jarlskog (World Scientific, Singapore, 1989), p. 175; Y. Nir and H.R. Quinn, Annu. Rev. Nucl. Part. Sci. **42**, 211 (1992).
- [13] For recent works, see, e.g., A.I. Sanda and Z.Z. Xing, Phys. Rev. D **56**, 6866 (1997); Z.Z. Xing, Eur. Phys. J. C **4**, 283 (1998); hep-ph/9712324; L. Randall and S. Su, hep-ph/9807377; and references therein.
- [14] See, e.g., L. Maiani, Report No. LNF-92/043(P); and references therein.




Figure 1: Illustrative plot for the time distribution of $\mathcal{A}(t)$ with $r \equiv \text{Re}\delta_B/\text{Im}\delta_B = \pm 1$ or ± 100 , where $x = 0.7$ and $y = 0.01$ have typically been taken.